



ORIGINAL ARTICLE

Numerical solution for Fredholm–Volterra integral equation of the second kind by using collocation and Galerkin methods

F.A. Hendi ^{a,*}, A.M. Albugami ^b

^a Department of Mathematics, Girls College of Education, King Abdul Aziz University, Jeddah, Saudi Arabia

^b Department of Mathematics, Girls College of Education, UAU, Makkah, Saudi Arabia

Received 11 March 2008; accepted 12 December 2009

Available online 23 December 2009

KEYWORDS

Integral equation;
Continuous kernel;
Collocation method;
Galerkin method

AMS

45E10
65R10

Abstract The Fredholm–Volterra integral equation of the second kind with continuous kernels with respect to position and time, is solved numerically, using the Collocation and Galerkin methods. Also the error, in each case, is estimated.

© 2009 King Saud University. All rights reserved.

1. Introduction

Many problems of mathematical physics, engineering and contact problems in the theory of elasticity lead to integral equations. The following references Muskhelishvili et al. (1953), Green (1969), Atkinson et al. (1976) and Delves

and Mohamed (1985), contain many different methods to solve the integral equations analytically. At the same time the numerical methods take an important place in solving the integral equations numerically. The references Linz et al. (1985), Kanwal et al. (1996), Atkinson et al. (1997) and Abdou and Mohamed (2002) contain many different methods for solving the integral equations numerically. The discussion of the Fredholm–Volterra integral equations numerically and analytically can be found in the works of Abdou and co-workers, see (Schiavane and Constanda, 2002; Abdou et al., 2003; Abdou and Salama, 2004), when the Fredholm integral term is considered in position and Volterra integral term in time. In all work of Abdou in Fredholm–Volterra integral equation when the kernel of position is continuous have not been solved.

Therefore, in this paper, we consider the Fredholm–Volterra integral equations of the second kind with continuous kernels with respect to position and time. The

* Corresponding author.

E-mail addresses: FALHENDI@kau.edu.sa (F.A. Hendi), abeer101aa@yahoo.com (A.M. Albugami).

1018-3647 © 2009 King Saud University. All rights reserved. Peer-review under responsibility of King Saud University.
doi:10.1016/j.jksus.2009.12.006



existence and uniqueness of the solution, under certain conditions, will be proved in the space $L_2[a, b] \times C[0, T]$, $0 \leq t \leq T \leq \infty$.

A numerical method is used to represent the Fredholm–Volterra equation in the form of a linear system of Fredholm integral equations where the existence and uniqueness of the system are discussed. Also we used the Collocation and Galerkin methods to obtain a linear system of algebraic equations, which is also solved numerically. Moreover the error estimate, in each case, is calculating.

2. The existence and uniqueness of the solution

Consider the Fredholm–Volterra integral equation

$$\mu\phi(x, t) - \lambda \int_a^b k(x, y)\phi(y, t)dy - \lambda \int_0^t F(t, \tau)\phi(x, \tau)d\tau = f(x, t) \quad (2.1)$$

where $k(x, y)$ and $F(t, \tau)$ are continuous functions which represent the kernel of Fredholm and Volterra integral terms, respectively. The known function $f(x, t)$ is called the free term of the integral equation, while $\phi(x, t)$ is unknown and called the potential function. Here the Fredholm is considered in position, while Volterra in time. The constant μ defines the kind of integral equation while the constant λ , may be complex, has a physical meaning.

In order to guarantee the existence of a unique solution of (2.1). We assume the following:

- (i) The kernel of position satisfies $|k(x, y)| \leq N_1$ for all, $a \leq x, y \leq b$, where N_1 is a constant.
- (ii) The positive continuous function $F(t, \tau) \in C([0, T] \times [0, T])$ for all $0 \leq t, \tau \leq T < \infty$ and satisfies $|F(t, \tau)| \leq N_2$ where N_2 is a constant.
- (iii) The given function and its norm is defined as: $f(x, t) \in L_2[a, b] \times C[0, T]$,

$$\|f(x, t)\| = \max_{0 \leq t \leq T} \int_0^t \int_a^b \{f^2(x, \tau)\}^{\frac{1}{2}} dx d\tau = N_3$$

where N_3 is a constant.

- (iv) The unknown function $\phi(x, t)$ satisfy the Lipschitz condition with respect to position $|\phi(x_1, t) - \phi(x_2, t)| \leq A(t)|x_1 - x_2|$ and Hölder condition with respect to time $|\phi(x, t_1) - \phi(x, t_2)| \leq B(x)|t_1 - t_2|^\alpha$, $0 \leq \alpha \leq 1$, and its norm is defined as

$$\|\phi(x, t)\| = \max_{0 \leq t \leq T} \int_0^t \int_a^b \{\phi^2(x, \tau)\}^{\frac{1}{2}} dx d\tau = N_4$$

3. The system of Fredholm integral equations

For representing (2.1) as a system of Fredholm integral equations we use the following numerical method, see Delves and Mohamed (1985) and Atkinson et al. (1997).

Divide the interval $[0, T]$ as

$$0 = t_0 < t_1 < \dots < t_k < \dots < t_N = T, \quad i.e. t = t_k, \\ k = 0, 1, 2, \dots, N.$$

Then, using the quadrature formula, Te Volterra term in (2.1) becomes

$$\int_0^{t_k} \phi(x, \tau) F(t_k, \tau)d\tau = \sum_{j=0}^k u_j F(t_k, t_j)\phi(x, t_j) + O(\tilde{h}_i^{\tilde{p}+1}), \\ (\tilde{h}_k \rightarrow 0, \tilde{p} > 0) \quad (3.1)$$

where $\tilde{h}_k = \max_{0 \leq j \leq k} h_j$, $h_j = t_{j+1} - t_j$

The values of k and the constant \tilde{p} depend on the number of derivatives of $F(t, \tau)$, for all $\tau \in [0, T]$, w.r.t. t , and $u_0 = \frac{1}{2}h_0$, $u_k = \frac{1}{2}h_k$, $u_i = h_i$, ($i \neq 0, k$).

Using (3.1) in (2.1) after letting $t = t_k$, $k = 1, 2, \dots, N$, We have

$$\mu\phi(x, t_k) = f(x, t_k) + \lambda \int_a^b k(x, y)\phi(y, t_k)dy + \lambda \sum_{j=0}^k u_j F(t_k, t_j)\phi(x, t_k) \quad (3.2)$$

or,

$$\mu\phi_k(x) = f_k(x) + \lambda \int_a^b k(x, y)\phi_k(y)dy + \lambda \sum_{j=0}^k u_j F_{k,j}\phi_j(x)\phi_k(x) \\ = \phi(x, t_k), f_k(x) = f(x, t_k), F_{k,j} = F(t_k, t_j), \quad (3.3)$$

the formula (3.3) become

$$\mu_n\phi_n(x) = G_n(x) + \lambda \int_a^b k(x, y)\phi_n(y)dy \quad (3.4)$$

where $\mu_n = \mu - \lambda u_n F_{n,n}$; $\lambda_n = \lambda u_n F_{n,n}$,

$$G_n(x) = f_n(x) + \lambda \sum_{j=0}^{n-1} u_j F_{n,j}\phi_j(x), \quad n = 0, 1, \dots, N$$

The formula (3.4) represents a linear system of Fredholm integral equations of the second kind, where $\lambda u_n F_{n,n} \neq \mu$.

Now, we will solve the linear system (3.4) using the Collocation method and Galerkin method.

3.1. Collocation method

Collocation method is based on approximating the solution $\phi(x, t)$ by a partial sum:

$$S(x, t_i) = \sum_{k=1}^N c_k(t_i)\psi_k(x) \quad (3.5)$$

of N linearly independent functions $\psi_1(x), \psi_2(x), \dots, \psi_N(x)$ on the interval (a,b). Therefore we have

$$\mu S_i(x) - \lambda \int_a^b k(x, y)S_i(y)dy \\ \approx f(x, t_i) + \lambda \sum_{j=0}^{i-1} w_j F_{i,j} S_j(x) + \varepsilon(x, c_1(t), c_2(t), \dots, c_N(t) + R(\tilde{h}_i^{p+1}) \quad (3.6)$$

Of course, if the approximate solution (3.5) is to be substituted into (3.4) for $\phi(x, t)$, there will be an error $\varepsilon(x, c_1(t), c_2(t), \dots, c_N(t))$. This error depends on x, t and the way for which the coefficients $c_1(t), c_2(t), \dots, c_N(t)$ are chosen in the formula (3.6). Let $t = t_i$, $i = 0, 1, 2, \dots, N$. Then using the quadrature formula, we have

$$\mu_i S_i(x_m) - \lambda \int_a^b k(x_m, y)S_i(y)dy \\ \approx f(x_m, t_i) + \lambda \sum_{j=0}^{i-1} w_j F_{i,j} S_j(x_m), i, m = 0, 1, 2, \dots, N \quad (3.7)$$

For determining the coefficients $c_1(t_i), c_2(t_i), \dots, c_N(t_i)$ of the approximate solution $S_N(x_N)$, from (3.5), in terms of the given N linearly independent functions $\psi_1(x), \psi_2(x), \dots$,

$\psi_N(x)$, perform the integration, then substitute $x = x_1, x_2, \dots, x_N$ for which the error $\varepsilon(x, c_1(t), c_2(t), \dots, c_N(t))$ vanishes.

Substituting from (3.5) in (3.6), we get:

$$\begin{aligned} & \mu_i \sum_{k=1}^N c_k(t_i) \psi_k(x_m) - \lambda \sum_{k=1}^N c_k(t_i) \int_a^b k(x_m, y) \psi_k(y) dy \\ & = f_{mi} + \lambda \sum_{j=0}^{i-1} \sum_{k=1}^N w_j F_{ij} c_k(t_j) \psi_k(x_m) \end{aligned} \quad (3.8)$$

3.2. Galerkin method

This method establishes the N conditions necessary for the determination of the N coefficients in Eq. (3.5):

By making the error $\varepsilon(x, c_1(t), c_2(t), \dots, c_N(t))$ in (3.6) orthogonal to N given linearly independent functions $\psi_1(x), \psi_2(x), \dots, \psi_N(x)$ on the interval (a, b) , i.e.

$$\int_a^b \psi_j(x) \varepsilon(x, c_1(t), c_2(t), \dots, c_N(t)) dx = 0 \quad (3.9)$$

Then from (3.6), we have

$$\begin{aligned} & \int_a^b \psi_j(x) [\mu_i S_i(x_m) - f(x_m, y) - \lambda \int_a^b k(x_m, y) S_i(y) dy \\ & - \lambda \sum_{j=0}^{i-1} w_j F_{ij} S_j(x) - R(\tilde{h}_i^{p+1})] dx = 0 \end{aligned} \quad (3.10)$$

Eq. (3.10) can be written in the form:

$$\begin{aligned} & \int_a^b \psi_j(x) [\mu_i S_i(x_m) - \lambda \int_a^b [k(x_m, y) S_i(y) dy \\ & - \lambda \sum_{j=0}^{i-1} w_j F_{ij} S_j(x_m) - R(\tilde{h}_i^{p+1})] dx \\ & = \int_a^b \psi_j(x) : f(x_m, t_i) dx \quad (i, m = 0, 1, \dots, N), \end{aligned} \quad (3.11)$$

where $R(\tilde{h}_i^{p+1})$ is the error from dividing the time and $\tilde{h}_i = \max_{0 \leq j \leq i} h_j, h_j = t_{j+1} - t_j$. The values of i and the constant p depend on the derivatives of $F(t, \tau)$, for all $\tau \in [0, T]$, with respect to t . Substituting from (3.5) into (3.11) we get

$$\begin{aligned} & \int_a^b \psi_j(x) [\mu_i \sum_{k=1}^N c_k(t_i) \psi_k(x_m) - \lambda \int_a^b k(x_m, y) \\ & \cdot \sum_{k=1}^N c_k(t_i) \psi_k(y) dy - \lambda \sum_{j=0}^{i-1} w_j F_{ij} \sum_{k=1}^N c_k(t_j) \psi_k(x_m)] dx \\ & = \int_a^b \psi_j(x) f(x_m, t_i) dx \end{aligned} \quad (3.12)$$

4. Examples

Example 4.1. Consider the integral equation:

$$\phi(x, t) = f(x, t) + \lambda \int_0^t \tau^2 \phi(x, \tau) d\tau + \lambda \int_a^b e^{x+y} \phi(y, t) dy$$

where the exact solution $\phi(x, t) = t^2 e^x$.

4.1. Using collocation method

In Eq. (4.1) we shall take $N = 2, a = 0, b = 1,$

$$f(x, t) = \frac{3}{2} t^2 e^x - \frac{1}{5} e^x t^5 - \frac{1}{2} t^2 e^x e^2$$

Let the approximate solution has the form of Eq. (3.5), the three independent functions are $\psi_1(x) = 1, \psi_2(x) = x, \psi_3(x) = x^2$. Substituting these values in Eq. (3.8), then solving the equations formulas when $x = 0, 0.5, 1.0,$ in this case $R = 0$. We get:

$$\begin{aligned} c_1(t_0) &= 0, \quad c_2(t_0) = 0, \quad c_3(t_0) = 0. \\ c_1(t_1) &= 0.0005130967957, \quad c_2(t_1) = 0.000432839493, \\ c_3(t_1) &= 0.000415594769. \end{aligned} \quad (4.1)$$

$$\begin{aligned} c_1(t_2) &= 0.002104378751, \quad c_2(t_2) = 0.001731361786, \\ c_3(t_2) &= 0.001662382707. \end{aligned} \quad (4.2)$$

So, the solution, for $t \in [0, 0.03]$, takes the form (see Table 1):

$$\begin{aligned} S(x, t_0) &= 0 \\ S(x, t_1) &= 0.0005130967957 + 0.000432839493x \\ &\quad - 0.000415594769x^2 \\ S(x, t_2) &= 0.002104378751 + 0.001731361786x \\ &\quad + 0.001662382707x^2 \end{aligned} \quad (4.3)$$

4.2. Using Galerkin method

As in collocation method, using (3.9) in (3.12), Choose three independent functions $\psi_1(x) = 1, \psi_2(x) = x, \psi_3(x) = x^2$ and three points $x = 0, 0.5, 1.0,$ when we assume $t \in [0, 0.03]$, then we have:

$$\begin{aligned} c_1(t_0) &= 0, \quad c_2(t_0) = 0, \quad c_3(t_0) = 0. \\ c_1(t_1) &= 0.00051949274, \quad c_2(t_1) = 0.00042025907, \\ c_3(t_1) &= 0.00041436290. \\ c_1(t_2) &= 0.00212991168, \quad c_2(t_2) = 0.0016810402, \\ c_3(t_2) &= 0.0016574556. \end{aligned} \quad (4.4)$$

Table 1 Values of the error E^C, E^G using collocation and Galerkin methods.

x	$\phi(x, t)$	E^C	E^G
$t = 0$			
0	0	0	0
0.5	0	0	0
1.00	0	0	0
$t = 0.0150000000$			
0	0.000225	0.0002880967957	0.00029449274
0.5	0.0003709622860	0.0004624529484	0.0004622507140
1.00	0.0006116134113	0.0007499176467	0.0007425012987
$t = 0.0300000000$			
0	0.0009	0.001204378714	0.00122991170
0.5	0.001483849144	0.001901806122	0.001900946656
1.00	0.002446453645	0.003051669518	0.003021953955

$$\begin{aligned}
 S(x, t_0) &= 0, \\
 S(x, t_1) &= 0.00051949274 + 0.00042025907x \\
 &\quad + 0.00041436290x^2, \\
 S(x, t_2) &= 0.00212991168 + 0.0016810402x \\
 &\quad + 0.0016574556x^2.
 \end{aligned}
 \tag{4.5}$$

So, the solution is taken from:

Example 4.2. Consider the integral equation:

$$\phi(x, t) = f(x, t) + \lambda \int_0^t \tau \phi(x, \tau) d\tau + \lambda \int_a^b e^{-y} \phi(y, t) dy, \tag{4.6}$$

where the exact solution $\phi(x, t) = te^{-x}$.

4.3. Using collocation method

In Eq. (4.6) we shall take $N = 2, a = 0, b = 1,$

$$f(x, t) = -0.00432t + e^{-x}t - 0.333e^{-x}t^4.$$

Let the approximate solution in the form of Eq. (3.5), then choose three independent functions $\psi_1(x) = 1, \psi_2(x) = x, \psi_3(x) = x^2$. Substituting these values in Eq. (3.8), then solving the equations formulas, when $x = 0, 0.5, 1.00,$ in this case $R = 0,$ we get:

$$\begin{aligned}
 c_1(t_0) &= 0, \quad c_2(t_0) = 0, \quad c_3(t_0) = 0. \\
 c_1(t_1) &= 0.008541939634, \quad c_2(t_1) = -.01412633660, \\
 c_3(t_1) &= 0.004644538564. \quad c_1(t_2) = 0.01708389624, \\
 c_2(t_2) &= -0.02825268914, \quad c_3(t_2) = 0.009289082405
 \end{aligned}
 \tag{4.7}$$

So, the solution, for $t \in [0, 0.03]$ takes the form:

$$\begin{aligned}
 S(x, t_0) &= 0, \\
 S(x, t_1) &= 0.008541939634 - 0.01412633660x + 0.004644538564x^2, \\
 S(x, t_2) &= 0.01708389624 - 0.02825268914x + 0.009289082405x^2.
 \end{aligned}
 \tag{4.8}$$

4.4. Using Galerkin method

We choose three independent functions $\psi_1(x) = 1, \psi_2(x) = x, \psi_3(x) = x^2$ and three points $x = 0, 0.5, 1.00,$ when we assume $t \in [0, 0.03],$ we have:

$$\begin{aligned}
 c_1(t_0) &= 0, \quad c_2(t_0) = 0, \quad c_3(t_0) = 0. \\
 c_1(t_1) &= 0.008459192683, \quad c_2(t_1) = -0.01395821152, \\
 c_3(t_1) &= 0.00463077286. \\
 c_1(t_2) &= 0.01691840427, \quad c_2(t_2) = -0.02791644232, \\
 c_3(t_2) &= 0.00926155092.
 \end{aligned}
 \tag{4.9}$$

So, the solution is taken the form (see Table 2):

$$\begin{aligned}
 S(x, t_0) &= 0, \\
 S(x, t_1) &= 0.008459192683 - 0.01395821152x \\
 &\quad + 0.00463077286x^2, \\
 S(x, t_2) &= 0.01691840427 - 0.02791644232x \\
 &\quad + 0.00926155092x^2.
 \end{aligned}
 \tag{4.10}$$

Table 2 Values of the error E^C, E^G using collocation and Galerkin methods.

x	$\phi(x, t)$	E^C	E^G
$t = 0$			
0	0	0	0
0.5	0	0	0
1.00	0	0	0
$t = 0.01500000000$			
0	0.015	0.006458060366	0.006540807317
0.5	0.009097959896	0.006458053921	0.006460179758
1.00	0.005518191618	0.006458050020	0.006386437595
$t = 0.03000000000$			
0	0.03	0.01291610376	0.01308159573
0.5	0.01819591979	0.01291609752	0.01292034895
1.00	0.01103638324	0.01291609374	0.01277287037

Acknowledgements

The authors would like to thank professor M.A. Abdou (University of Alexandria, Egypt) for his helpful remarks and suggestions.

References

Abdou, M.A., Mohamed, K.I., Ismail, A.S., 2002. Toeplitz Matrix and Product Nystrom Methods for Solving the Singular Integral Equation, *Le Mathematical*, vol. Lv 11, Fasc 2, pp. 21–37.

Abdou, M.A., Salama, F.A., 2004. Volterra – Fredholm integral equation of the first kind and spectral relation ships. *J. Appl. Math. Comput.* 153, 141–153.

Abdou, M.A., Mohamed, K.I., Ismail, A.S., 2003. On the numerical solutions of Fredholm–Volterra integral equation. *J. Appl. Math. Comput.* 146, 713–728.

Atkinson, K.E., 1976. *A Survey of Numerical Methods for the Solution of Fredholm Integral Equation of the Second Kind*. Philadelphia.

Atkinson, K.E., 1997. *The Numerical Solution of Integral Equation of the Second Kind*. Cambridge.

Delves, L.M., Mohamed, J.L., 1985. *Computational Methods for Integral 8 Equation*. Cambridge.

Green, C.D., 1969. *Integral Equation Methods*. Nelson, NewYork.

Kanwal, R.P., 1996. *Linear Integral Equations Theory and Technique*. Boston.

Linz, P., 1985. *Analytical and Numerical Methods for Volterra Equations*. SIAM, Philadelphia.

Muskhelishvili, N.I., 1953. *Singular Integral Equations*. Noordhoff, Groningen, The Netherlands.

Schiavane, P., Constanda, C., Mioduchowski, A., 2002. *Integral Methods in Science and Engineering*. Birkhauser, Boston.