

Seminoma of an Undescended Testis Presenting As Incomplete Large Bowel Obstruction

**ADNAN A. MERDAD, FRCS(C),
HISHAM A. MOSLI, FRCS(C), and OSSAMA I. NASSIF, FRCP(C)**
*Associate Professor, Department of Surgery; Associate Professor,
Department of Urology; and Assistant Professor, Department of Pathology,
Departments of Surgery, Urology and Pathology, King Abdulaziz University
Hospital, Jeddah, Saudi Arabia*

ABSTRACT. Seminoma is the most common neoplasm to arise in the undescended testicle. The clinical presentation of such a tumor is either as an abdominal mass or abdominal pain. A patient with a seminoma developing in an undescended testicle which presented as an incomplete large bowel obstruction is being reported. To our knowledge, such presentation has not been reported previously. The clinical presentation, investigations, treatment modalities, and long term follow-up result are described.

Introduction

The undescended testicles carry a higher potential for malignant transformation than the normally descended intrascrotal testes. This risk is highest among the intra-abdominal testes.^[1] The intra-abdominal germ cell tumors arising from an undescended testis may present as painless mass or acute abdominal pain secondary to torsion of the pedicle or hemorrhage within the tumor.^[2-5]

We are reporting a patient with a seminoma arising in an undescended testicle who presented in a state of incomplete large bowel obstruction.

Correspondence to: Dr. A. Merdad, Consultant Surgeon, King Abdulaziz University Hospital, Department of Surgery, P.O. Box 6615, Jeddah 21452, Saudi Arabia.

Case Report

A 22-year old male was referred to our unit with a chief complaint of progressive difficulty in passing stools which were frequently covered with bright red blood for the last 8 months. The patient was gradually losing weight. He had been taking laxatives regularly for the last two months. He also suffered left sided lower abdominal pain of moderate severity.

On examination, he looked moderately ill, pale but not jaundiced. His abdomen was slightly distended, soft, and with no palpable masses. He was slightly tender in his left lower abdominal quadrant. Examination of the genitalia revealed absence of the left testicle. Rectal examination revealed at 6 cm from anal verge a hard fungating, ulcerating lesion that bled easily on touch. The lesion involved the rectum circumferentially. A rigid sigmoidoscope could not pass beyond it.

Multiple biopsies were obtained. Histologically, the picture was of a malignant neoplasm compatible with a germ cell testicular tumor with features typical of seminoma (Fig. 1). Immunohistochemical studies (Avidin Biotin technique) showed that the cells infiltrating the stroma were positive for placental alkaline phosphatase, but negative for cytokeratin, epithelial membrane antigen, and leukocyte common antigen. Meanwhile, barium enema (Fig. 2a & b) and computerized tomographic





FIG.1. High power magnification of transrectal biopsy shows groups of large seminoma cells within the stroma which is heavily infiltrated by lymphocytes.



(a)



(b)

FIG.2a & b. Double contrast barium enema AP + lateral projections showing 10 cm long segment at the rectosigmoid with narrowing and irregularity.

examination of the abdomen (Fig. 3) showed a large retroperitoneal solid mass invading the rectum. The testicular tumor markers, alpha fetoprotein and serum human betachorionic gonadotrophin were within the normal limits as were the renal and liver function tests.




FIG.3. CT scanning image at the upper rectum showing thickening of the wall, irregularities and narrowing of the lumen.

The patient was treated with both radiotherapy and cyclic courses of platinum based combination chemotherapy of VAB (Vinblastine-Adriamycin-Bleomycin) protocol. The pulmonary function tests as well as the hemopoietic picture were carefully monitored during chemotherapy.

He has been followed up regularly every three to six months for the past three years. All his blood work is normal. Repeat CT scans of abdomen showed complete regression of the tumour (Fig. 4). His last sigmoidoscope two and half years after his treatment showed scarring at the area of the tumor with minimal narrowing and multiple biopsies showed no evidence of recurrence.

Discussion

Seminoma is the most common single histologic type of all germinal testicular neoplasms.^[6] Primary germ cell tumours originating from undescended testes appear to carry no worse prognosis than those arising in the scrotal testis.^[7]



FIG. 4. CT scanning image at the same region as Fig. 2 post chemoradiotherapy showing marked improvement in the rectum. The lumen is still narrowed with some wall thickening.

Approximately 25% of seminoma patients have metastatic disease at their initial presentation.^[8] Until recently, because of its high radiosensitivity, seminoma has been primarily treated with irradiation in all stages. Patients presenting with a retroperitoneal tumor > 5 cm in diameter, supradiaphragmatic and extralymphatic metastases are labelled to have an advanced disease. This latter group constitutes 15% of all seminoma patients. It represents an exception in regards to primary treatment since megavoltage irradiation has not proven adequate because of the relatively low cure rate and the high relapse rate.^[8] When treated with primary radiotherapy alone the cure rate for this group of patients is 65% only, this is considered unacceptably low when compared to the cure rate of 93-100% using the same modality for early stages of seminoma. Systemic chemotherapy has become the logical therapeutic alternative and supplement to radiotherapy and the preferred initial treatment of seminoma patients presenting with advanced disease.

Patients with advanced disease initially treated with platinum based chemotherapy programs and irradiation and/or salvage chemotherapy for residual or recurrent tumors can expect an overall cure rate of 88-92%.^[9]

The reported patient three years after his treatment is back to his normal activities and shows no evidence of recurrence.

References

- [1] **Witaker RH.** Management of the undescended testis. *Br J Hosp Med* 1970; **4**: 25-37.
- [2] **Munro AJ, Duncan W, Webb JN.** Extragonadal Presentations of Germ Cell Tumours. *Br J Urol* 1983; **55**(5): 547-554.
- [3] **Awad RM, Shetty SD, Ibrahim A.** Intra-abdominal seminoma presenting as an acute abdomen. *Ann Saudi Med* 1988; **8**(4): 290-291.
- [4] **Sarma DP, Weilbaecher TG, Hatem AA.** Seminoma arising in undescended testis clinically presenting as acute appendicitis. *J Surg Oncol* 1986; **31**(1): 44-47.
- [5] **Packham DA.** Torsion of an intra-abdominal seminoma of the testis. *Br J Surg* 1965; **52**: 25-27.
- [6] **Mostofi FK, Sisterhenn I, Davis CJ.** Evaluation of WHO classification with tumor markers in 1,000 testicular tumors. *J Urol* 1984; **131**(4): 195 A.
- [7] **Hogan P, Smedley H, Sikora K.** Abdominal pain as a presenting symptom of male germ cell tumour. *Br J Urol* 1985; **57**(2): 197-199.
- [8] **Wettlaufer JN.** The Role of Radiotherapy and Chemotherapy in Advanced Seminoma, in: *EORTC Genitourinary Group Monograph: Progress and Controversies in Oncological Urology II*. New York: Alan R. Liss, Inc, 1988: 395-404.
- [9] **Wettlaufer JN.** The management of advanced seminoma. *Sem Urol* 1984; **II**(4): 257-263.

ورم منوي خصوي في خصية غير نازلة يظهر على شكل انسداد غير كامل بالأمعاء الغليظة

عدنان عبد المعطي مرداد و هشام أحمد موصلي و أسامة إبراهيم ناصف
 أستاذ مشارك ، قسم الجراحة أستاذ مشارك ، قسم المسالك البولية أستاذ مساعد بقسم علم الأمراض
 مستشفى جامعة الملك عبد العزيز ، جدة - المملكة العربية السعودية

المستخلص . إن الورم المنوي الخصوي هو أكثر الأورام حدوثاً في الخصية المتحجرة (غير النازلة) . سريريًا يظهر هذا الورم على شكل ألم أو كتلة بالبطن .

هنا ، تُسجل حالة مريض بورم منوي خصوي تكونت في خصية متحجرة وأدت إلى انسداد غير كامل بالأمعاء الغليظة .

على حد علم المؤلفين هذه أول حالة تسجل على هذا النحو . حالة المريض وفحوصاته وطريقة علاجه ومتابعته سوف تشرح بقدر من التفصيل . *

